

RESEARCH

Open Access



Depression and health-related quality of life of patients with type 2 diabetes attending tertiary level hospitals in Dhaka, Bangladesh

Manish K. Namdeo^{1*} , Sarita Verma² , Rajat Das Gupta³, Rubana Islam⁴ , Shaila Nazneen⁵ and Lal B. Rawal⁶

Abstract

Introduction Type 2 diabetes mellitus (T2DM) and depression are closely linked. People with T2DM are at increased risk of developing depression and vice versa. T2DM and depression comorbid conditions adversely affect Health-Related Quality of Life (HRQOL) and management of T2DM. In this study, we assessed depression and HRQOL among patients with T2DM in Dhaka, Bangladesh.

Methods A cross-sectional study was conducted in two tertiary-level hospitals in Dhaka, Bangladesh. Data were collected from 318 patients with T2DM. A set of standard tools, PHQ-9 (for assessing depression) and EuroQol-5D-5L (for assessing the HRQOL), were used. Statistical analyses, including Chi-square and Fisher's exact tests, Wilcoxon (Mann-Whitney), and Spearman's correlation coefficient tests, were performed using SPSS (v.20).

Results The majority of the patients (58%) were females, with a mean age (standard deviation) of 52 ± 10 years, and 74% of patients lived in urban areas. The prevalence of depression was 62% (PHQ-9 score ≥ 5). Over three-quarters (76%) reported problems in the anxiety/ depression dimension of EQ-5D, followed by pain/discomfort (74%), mobility (40%), self-care (36%), and usual activities (33%). The depression and T2DM comorbid condition were associated with all the five dimensions of EQ-5D (χ^2 statistics with $df = 1$ was 52.33, 51.13, 52.67, 21.61, 7.92 for mobility, self-care, usual activities, pain/discomfort, and anxiety/ depression dimensions respectively, $p < 0.01$). The mean EQ-5D index (0.53 vs. 0.75) and the mean EQ-5D VAS (65 vs. 76) both showed lower values in T2DM patients with depression compared to T2DM patients without depression (Wilcoxon test, $p < 0.001$).

Conclusions We conclude that the majority of the patients with T2DM had comorbid conditions, and the HRQOL was negatively affected by comorbid depression in T2DM patients. This suggests the importance of timely screening, diagnosis, treatment, and follow-up of comorbid depression in T2DM patients to improve overall health and QOL.

Keywords Depression, HRQOL, Comorbidity, Bangladesh, Type 2 diabetes

*Correspondence:
Manish K. Namdeo
drmanish1@gmail.com
Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Background

Type 2 diabetes mellitus (T2DM) is a major public health problem worldwide. Nearly half a billion people are living with T2DM worldwide, of whom 75% live in low- and middle-income countries (LMICs) [1]. The situation is particularly alarming in South-East Asia, where 87.6 million people were affected with diabetes in 2019, resulting in 1.2 million attributable deaths to T2DM [2]. More specifically, Bangladesh follows along with these estimations [3, 4] amidst the aggravating background of a health system where communicable diseases are still prevalent [5]. Bangladesh is among the top 10 countries with the most cases of undiagnosed T2DM in adults [2]. The burden of T2DM involves high costs to patients (US\$88 a year) and providers (US\$54–64 a year) in Bangladesh [2, 6]. These costs include medical advice, investigations, treatment, travel, productivity loss, and carers' expenses. As reported by the IDF, the annual health expenditure per person with T2DM in Bangladesh during 2019 was estimated at US\$64. This figure represents the expenses directly linked to diabetes, irrespective of who pays, whether patients, private or public payers, or the government [2].

Evidence shows that people with T2DM are 2–3 times more likely to develop depression than those without diabetes, and vice versa [7, 8]. In Bangladesh, depression is a major issue, affecting 41% of the general population and 34% of individuals with chronic medical conditions. A study among university students in Bangladesh reported that 65% of patients exhibited depressive symptoms [9]. The co-occurrence of multiple chronic illnesses, such as diabetes and depression, demonstrates a strong interconnection [10], with the odds of depression doubling in T2DM patients [11–13]. Two hypotheses explain this relationship. The first hypothesis suggests that depression precedes diabetes, as changes in counter-regulatory hormones, glucose transport mechanisms, and immune-inflammatory responses contribute to the development of depression [12]. These alterations lead to disruptions in glucose metabolism, resulting in B-cell dysfunction and insulin resistance [12]. The second hypothesis proposes that the chronic nature of diseases like diabetes, along with associated complications, induces stress that can trigger depressive illness in diabetic patients [14]. The relationship between depression and diabetes is bidirectional, and this supports both hypotheses; hence, the validity of one hypothesis does not negate the possibility of the other hypothesis being true [13, 15].

Complications and multi-organ involvement in T2DM, along with high treatment costs, impact the Quality of Life (QoL) of patients and those around them [16]. QoL is an individual's perception of their life within the context of their culture and value systems, considering their

goals, expectations, standards, and concerns [17]. QoL encompasses a broader concept of how diseases hinder one's ability to fulfill normal roles, reflected as self-perceived Health-Related Quality of Life (HRQOL) in terms of physical and mental well-being [16, 18, 19].

T2DM itself reduces mean HRQOL scores compared to healthy individuals [20, 21]. The coexistence of depression and diabetes worsens HRQOL, leading to medication non-compliance, increased complications, and elevated suicide risk [21–25]. Assessing HRQOL is valuable in understanding the psychosocial impact of T2DM and coexisting conditions [26–28]. Understanding the psychological dimensions of diseases can lead to improved treatment approaches [29]. Research findings indicate that the incorporation of mental health services into chronic disease care has shown positive outcomes, leading to improvements in both physical health conditions and depression [30, 31].

Studies in Bangladesh on HRQOL have been conducted in various groups, such as adolescents with cerebral palsy [32], leprosy patients [33], older people [34], postpartum mothers [35], T2DM patients [36, 37], and migrant workers [38], but research on T2DM patients with comorbid depression remains unassessed. This study offers unique insights, as it is the first documented study using both PHQ-9 and EQ-5D to assess depression comorbidity in T2DM patients. Considering the ongoing COVID-19 pandemic, mental health support for individuals with diabetes has become even more critical [39]. This further emphasizes the importance of publishing such studies. In this study, we determined the general characteristics and health status (body mass index (BMI), blood pressure, comorbid depression, and HRQOL) among patients with T2DM. We also examined the association between depression and HRQOL.

Methods

Study design and setting

This cross-sectional study was conducted between November 2014 and January 2015. A total of 318 patients with T2DM were enrolled in the study at two medical centers in Dhaka, Bangladesh. The first one, BIHS, is a medical teaching institute under the Health Care Development Project (HCDP) of Bangladesh Diabetic Society, attached to a 250-bed diabetic tertiary care hospital [40], but also linked to smaller regional hospitals and ten urban healthcare centers in and around Dhaka [40]. The second one, DMCH, is a medical college attached to a 2,300-bed hospital [41]. Its Department of Medicine provides outpatient services to diabetic patients on Mondays and Saturdays and is attended by patients from all over the country [41].

Participants

The study population comprised patients with clinically diagnosed T2DM. The inclusion criteria for the study participants were confirmed cases of T2DM, aged between 30 to 70 years, attending the outpatient departments (OPDs) of the BIHS hospital and DMCH during the study period, and being willing to participate in the study. Pregnant women were excluded, as pregnancy itself may affect the HRQOL in a woman [42]. Similarly, severely ill patients who required emergency care and could not respond to survey questions were excluded.

Sampling and sample size

Patients attending the OPDs of BIHS and DMCH during the study period were our potential respondents and were conveniently sampled. Patients waiting for their turn for consultation or collecting the results of the routine investigation in the waiting area were requested to participate in the study. 318 T2DM patients (280 from BIHS and 38 from DMCH) were interviewed from the study sites.

The minimum required sample size was calculated using the formula $n = Z^2 pq/d^2$ (Where $z = 1.96$, $p =$ the proportion of T2DM patients in Bangladesh who reported problems in anxiety/depression dimension of EuroQol-5D (EQ-5D), i.e., 73.6%, $q = 1 - p$, and $d =$ the allowable error of known prevalence, i.e., 5%) [36]. The present study estimated the sample size based on the prevalence of anxiety/depression in EQ-5D-5L dimensions because this was the most relevant dimension associated with the study objective [43]. With a 95% confidence interval and a 5% error, the calculated sample size was 299 [44]. Additional participants were included in the sample size calculation to account for any missing

data, resulting in 318 patients being interviewed for the study.

Study tools

We administered a survey questionnaire written in the Bangla language, comprising four sections: (I) socio-demographic characteristics; (II) physical measurements; (III) EQ-5D-5L for health-related quality of life; and (IV) PHQ-9 (Patient Health Questionnaire-9) for assessing depression (Table 1).

To measure the health-related quality of life (HRQOL), we used the standardized EuroQol-5D-5L (EQ-5D-5L) questionnaire. When evaluating HRQOL, different instrument types can be used, including disease-specific, generic, and utility instruments. The EQ-5D is considered the most appropriate generic instrument because of its broad applicability across the population, facilitating HRQOL comparisons [45]. Using EQ-5D provides a simpler, more cost-effective, and comprehensive evaluation of HRQOL, making it valuable for assessing health outcomes in diverse patient populations and conditions. The EQ-5D captures aspects of health, such as mobility, self-care, usual activities, pain/discomfort, and anxiety/depression. It comprises three measurements: (1) the EQ-5D descriptive system, which shows the distribution of individuals across five levels in each dimension; (2) the Visual Analogue Scale (VAS), a continuous value representing general health status; and (3) the EQ-5D index, which assigns a value to each EQ-5D health state, yielding a single composite index [46, 47]. To assess depression levels, we utilized a previously validated Bengali version of the PHQ-9, available for free download on the PHQ website [48, 49]. The PHQ-9 consists of nine items on a 4-point Likert scale, with standard cut-off scores

Table 1 Study variables and tools

Sections	Variable	Type of Variable	Measurement tools	Assessed information
Section I	Socio-demographic characteristics	Independent	Questionnaire	Age, sex, religion, marital status, education, occupation, income, expenditure, household members, area of residence, age at diagnosis of the illness, and years since the diagnosis
Section II	Physical measurements	Independent	Secondary data from patients' record	Height, weight, Blood pressure
Section III	Comorbid depression	Independent	PHQ 9	A. PHQ 9 score B. Severity of depression: 1. No; 2. Mild; 3. Moderate; 4. Moderately severe; 5. Severe C. Presence of depression 1. Diabetes with no depression 2. Diabetes with depression
Section IV	HRQOL	Dependent	EQ-5D-5L	A). EQ-5D-5L descriptive system dichotomized for all the five dimensions into the problem and no problem B). Visual analogue scale (VAS) C). EQ-5D-5L index value

used to classify symptoms of depression as minimal/none (0–4), mild (5–9), and moderate to severe (≥ 10) [50].

Data collection and processing

We pre-tested the survey questionnaire among a similar population at BIHS Hospital prior to data collection. Data was collected through face-to-face interviews conducted by key researchers and three trained assistants. Each interviewer was assigned a unique code (e.g., K011, S012), and the questionnaires were numbered serially. Patients were identified by combining the interviewers' code and questionnaire serial number (e.g., K011-001, S012-199). Each interview lasted 25–30 minutes, with approximately 7–8 sets of questionnaires completed by each interviewer daily, totaling 35–40 interviews. Questionnaires were checked daily for accuracy and completeness, and feedback was provided to interviewers regularly.

Cronbach's alpha for the PHQ-9 scale was 0.824, and the correlations between the nine items of the PHQ-9 and total PHQ-9 scores were significant at the 0.01 level, ranging from 0.610 to 0.769. Likewise, Cronbach's alpha for the EQ-5D scale was 0.816. The correlations between five items of the EQ-5D and total EQ-5D scores were

significant at the 0.01 level, ranging from -0.602 to -0.794 .

Data analysis

The data were analyzed using SPSS (v.20) [51]. Outliers in continuous variables were identified using Tukey's fence test, and extreme outliers were described using median and interquartile ranges. Categorical variables were analyzed using frequency tables, Chi-square, and Fisher's exact tests. The EQ-5D index and EQ-VAS (dependent variables) were found to have non-normal distributions with negative skewness in the Shapiro-Wilk test ($p < 0.05$). The histogram of the EQ-5D VAS revealed multiple spikes corresponding to numbers ending with 0 or 5, indicating a preference for these digits among participants (Figs. 1 and 2). Similar non-normal distributions have been observed in previous studies, warranting the use of non-parametric tests for comparing means [52].

The PHQ-9 questionnaire comprises nine items rated on a 4-point Likert scale. Standard cut-off scores were used to categorize depression as minimal/none (0–4), mild (5–9), and moderate to severe (≥ 10) [50]. In this study, the scores were further collapsed into two categories: no depression (< 5) and depression (≥ 5) [8]. This




Fig. 1 Histogram EQ-5D index: The figure represents a histogram displaying the distribution of EQ-5D index scores in a population. The EQ-5D index is a measure of health-related quality of life, ranging from 0 (representing death) to 1 (representing perfect health). The x-axis of the histogram shows the range of EQ-5D index scores, with the minimum score of 0 on the left and the maximum score of 1 on the right. The y-axis shows the frequency of individuals with each EQ-5D index score




Fig. 2 Histogram EQ-5D VAS: The histogram in Figure represents the distribution of EQ-5D Visual Analog Scale (VAS) scores among the study population. The EQ-5D VAS is a measure of health-related quality of life, ranging from 0 to 100, where a score of 100 represents perfect health and 0 represents worst imaginable health state. The x-axis shows the range of EQ-5D VAS scores, and the y-axis represents the frequency of individuals with each score. The histogram is divided into several bars, each representing a range of EQ-5D VAS scores

categorization created two groups among the patients: T2DM patients with depression and T2DM patients without depression. The EQ-5D dimensions were categorized as no problems (level 1) and problems (levels 2 to 5) [53]. Although dichotomizing the EQ-5D dimensions may result in oversimplification of the assessment, it can be useful in standardizing assessments across diverse patients and settings. Additionally, it offers a standardized approach for analysis and enables statistical comparisons. Chi-square, Fisher's exact test, and Wilcoxon (Mann-Whitney) tests were used to examine associations and differences between variables. Significance was set at $p < 0.05$. Spearman's correlation coefficient assessed relationships between patient characteristics, PHQ-9 scores, and the EQ-5D index. A trend plot was developed to illustrate the relationship between depression levels and the HRQOL index.

Results

Socio-demographic characteristics of the patients

Among the 318 enrolled patients, the majority ($n = 184$, 58%) were female, and most ($n = 280$, 88%) were from the BIHS hospital. About three-quarters ($n = 234$, 74%)

of the patients were from an urban area. Most of the patients ($n = 303$, 95%) were married and Muslims ($n = 312$, 98%) by religion. Most of the patients were educated, and 12% ($n = 38$) had no formal education. More than half ($n = 176$, 55%) of the patients were homemakers. Their median monthly household income was 321.50 USD (interquartile range (IQR) = 450.10 USD). After excluding one extreme outlier with an income of 15432.06 USD per month, the average monthly household income of the patients ($n = 317$) was 492.36 USD (standard deviation (SD) = 565.89 USD). The average monthly household expenditure of the patients was 407.19 USD (SD = 392.95 USD). The reported pattern showed that the majority of the patients had income (46%, $n = 146$) and expenditure (49%, $n = 155$) in the 10,000–30,000 BDT (128.60–385.80 USD) category. The mean age of the patients was 52 (SD = 11) years (median = 52 years). The majority ($n = 179$, 56%) were in the 41–60 age group. The median age at diagnosis of diabetes was 44 years (IQR = 13) (mean age at diagnosis 44; SD = 10 years). More than half ($n = 179$; 56.29%) of the patients had a family history of T2DM.

Anthropometric and biomedical measurements

The mean body mass index (BMI) of the patients ($n = 259$) was 20.18 (SD = 3.03) kg/meter². The mean systolic blood pressure of the patients ($n = 284$) was 124.95 (SD = 15.37) mmHg (median = 120; IQR = 10 mmHg), and mean diastolic blood pressure was 78.86 (SD = 7.99) mmHg (median = 80; IQR = 05 mmHg).

PHQ-9 for assessing depression

The proportion of depression among T2DM patients was 62% (PHQ-9 score ≥ 5). Among all the patients ($N = 318$), 38% ($n = 122$) had none/minimal depression (PHQ-9 score 0–4); 30% ($n = 94$) had mild depression (score 5–9); 23% ($n = 73$) had moderate depression (PHQ-9 score 10–14); 7% ($n = 22$) had moderately severe depression (PHQ-9 score 15–19); 2% ($n = 7$) had severe depression (PHQ-9 score 20–27) (Table 2).

EQ-5D for assessing health-related quality of life

About 40% of patients had problems in mobility, 36% had problems in self-care, 33% had problems in usual activities, 74% had problems in pain/discomfort, and 76% had problems in the anxiety/depression dimension of the EQ-5D (Table 3).

Table 2 Percentage of patients in different depression severity groups ($N = 318$)

Severity of depression (PHQ 9 score)	Count	N (%)
None (0–4)	122	38
Mild depression (5–9)	94	30
Moderate depression (10–14)	73	23
Moderately severe depression (15–19)	22	7
Severe depression (20–27)	7	2

Table 3 Frequency of reported problem by dimensions of HRQOL ($N = 318$)

Dimensions		Count	N%
Mobility	No problem	191	60
	Problem	127	40
Self-care	No problem	204	64
	Problem	114	36
Usual Activities	No problem	212	67
	Problem	106	33
Pain/Discomfort	No problem	84	26
	Problem	234	74
Anxiety/Depression	No problem	77	24
	Problem	241	76

EQ-5D index and EQ-5D VAS

The mean EQ-5D index was 0.62 (SD = 0.22) (median = .66, IQR = 0.24). Mean EQ-5D VAS was at 69 (SD = 20) (median = 70, IQR = 25). Within their respective categories, individuals in the younger age group, males, Hindus, those who have never been married, without a family history, belonging to smaller households (less than four family members), having a higher education level, working as laborers, with higher household income, shorter illness duration, without depression, residing in urban areas, and having normal systolic/diastolic blood pressure, exhibited higher EQ-5D index and VAS scores (Table 4).

Association of socio-demographic characteristics and comorbid depression with HRQOL

The proportion of patients having problems in all five dimensions of the EQ-5D (56 patients having problems in mobility vs. 15 patients with no problem, 51 with problems vs. 11 with no problem in self-care, 48 with problems vs. 9 with no problem in usual activity, 83 with problems vs. 59 no problem in pain discomfort, and 81 with problems vs. 61 no problem in anxiety/depression dimension) was significantly higher among those who showed the presence of depression (PHQ-9 score ≥ 5) (Table 5). The relationship between the EQ-5D dimensions and various demographic factors is presented in Table 5. The study revealed significant Spearman's correlations ($p < 0.01$) between the EQ-5D index and various demographic and health-related factors, including age ($r = -.21$), sex ($r = -.27$), duration of diagnosis ($r = -.12$), education ($r = .26$), and PHQ-9 severity score ($r = -.61$). Additionally, significant correlations ($p < 0.01$) were observed between the EQ-5D VAS and several factors, such as sex ($r = -.15$), monthly income ($r = .23$), expenditure ($r = .22$), BMI ($r = .15$), education ($r = .20$), and PHQ-9 severity score ($r = -.38$), as presented in Table 6.

Table 7 presents the average PHQ-9 scores, EQ-5D index, and VAS scores based on demographic characteristics and the presence of depression. Notably, our analysis revealed that individuals with depression had significantly lower EQ-5D index values compared to those without depression (mean EQ-5D index 0.53 vs. 0.75; median 0.54 vs. 0.72; Kruskal-Wallis test p -value < 0.05). Moreover, the EQ-5D VAS scores were significantly lower in individuals with depression (mean EQ-5D VAS 65 vs. 76; median 70 vs. 80; Kruskal-Wallis test p -value < 0.05). When exploring the severity of depression and its relationship with HRQOL, the study found that the mean EQ-5D index decreased with the increasing severity of depression (Fig. 3). The

Table 4 Mean EQ-5D index and mean EQ-5D VAS (N = 318)

		N (%)	EQ-5D-5L index Mean (SD)	EQ-5D-5L VAS Mean (SD)
Sex	Male	134(42)	.68(.19)	73(20)
	Female	184(58)	.57(.23)	67(19)
Age groups (years)	30–39	38(12)	.69(.21)	64(18)
	40–49	79(25)	.64(.19)	70(21)
	50–59	100(31)	.62(.19)	72(18)
	60 +	101(32)	.56(.25)	68(21)
Marital status	Never married	7(2)	.67(.25)	71(28)
	Married	303(95)	.61(.22)	69(20)
	Widow/Widower	7(2)	.69(.12)	79(11)
	Separated/Divorced	1(1)	.72	70
Family History of T2DM	No	139(44)	.64(.18)	68(19)
	Yes	179(56)	.60(.24)	70(20)
Household members	< 4 or = 4	145(46)	.64(.21)	70(20)
	> 4	173(54)	.60(.22)	69(20)
Education	No formal education	38(12)	.54(.26)	61(21)
	Primary education	103(32)	.57(.24)	66(21)
	Higher sec. education	108(34)	.64(.18)	72(18)
	College and above	69(22)	.70(.18)	75(20)
Occupation	Unemployed	3(1)	.56(.06)	65(22)
	Service	54(17)	.68(.20)	74(23)
	Business	32(10)	.66(.19)	72(23)
	Laborer	7(2)	.75(.22)	74(11)
	Farming	5(2)	.63(.15)	60(10)
	House maker	176(55)	.56(.22)	67(19)
	Retired	41(13)	.70(.17)	74(16)
Monthly household income (BDT)	< 10,000	52(16)	.53(.26)	63(20)
	10,000–30000	146(46)	.63(.21)	67(20)
	30,000–60000	83(26)	.62(.20)	73(18)
	60,000–90000	11(3)	.69(.17)	81(14)
	> 90,000	26(8)	.64(.19)	79(16)
Monthly household expenditure (BDT)	< 10,000	58(18)	.56(.25)	64(19)
	10,000–30000	155(49)	.63(.21)	68(20)
	30,000–60000	80(25)	.62(.21)	73(19)
	60,000–90000	9(3)	.72(.15)	83(16)
	> 90,000	16(5)	.62(.21)	80(18)
BMI	< 18.5 (Underweight)	9(3)	.62(.20)	66(20)
	18.5 to < 23 (Normal wt.)	83(26)	.62(.20)	72(20)
	23 to < 27.5 (Over wt.)	144(45)	.69(.20)	74(18)
	≥ 27.5 (Obese)	82(26)	.55(.27)	69(27)
Duration of diagnosis	Duration < = 3 years	108(34)	.66(.19)	70(18)
	Duration 3–6 years	53(17)	.62(.20)	69(21)
	Duration 6–9 years	47(14)	.59(.18)	69(18)
	Duration > 9 years	110(35)	.58(.25)	69(22)
Depressive symptoms	No depression (< 5 score)	122(38)	.75(.14)	76(16)
	Depression (≥ 5 score)	196(62)	.53(.22)	65(21)
Area of residence	Urban	234(74)	.63(.21)	71(20)
	Rural	39(12)	.49(.27)	60(21)
	Peri-Urban	45(14)	.67(.15)	70(18)

Table 4 (continued)

		N (%)	EQ-5D-5L index Mean (SD)	EQ-5D-5L VAS Mean (SD)
Systolic BP	Normal SBP	47(17)	.72(.16)	68(23)
	Prehypertension	189(67)	.63(.20)	71(18)
	Stage 1 Hypertension	29(10)	.49(.25)	67(25)
	Stage 2 Hypertension	19(7)	.57(.22)	66(19)
Diastolic BP	< 80 (Normal)	70(24.6)	.67(.18)	73(20)
	80–99 (Prehypertension)	168(59.2)	.62(.21)	69(20)
	90–99 (Stage 1 HTN)	35(12.3)	.57(.22)	73(18)
	> 100 (Stage 2 HTN)	11(3.9)	.63(.19)	50(18)

Table 5 EQ-5D dimensions by selected demographic characteristics (N = 318)

		N (%)	Mobility Problem%	Self-care Problem%	Usual activities Problem%	Pain/Discomfort Problem%	Anxiety/Depression Problem%
Sex	Male	134(42)	32	26	22	63	73
	Female	184(58)	46	43	42	82	78
χ^2 (df)			5.946(1)	9.533(1)	14.245(1)	14.151(1)	.887(1)
p-value			.015*	.002*	.000*	.000*	.346
Age (years)	30–39	38(12)	26	24	24	58	68
	40–49	79(25)	29	20	23	78	82
	50–59	100(31)	42	36	34	78	77
	60+	101(32)	51	52	45	71	72
χ^2 (df)			12.590(3)	22.942(3)	11.291(3)	7.064(3)	3.696(3)
p-value			.006*	.000*	.010*	.070	.296
Householder members	≤ 4	145(46)	32	34	32	72	74
	> 4	173(54)	46	38	34	75	77
χ^2 (df)			6.289	.490	.101	.188	.577
p-value			.012*	.484	.750	.665	.448
Education	No education	38(12)	55	55	53	76	71
	Primary edu	103(32)	50	38	37	83	81
	Higher sec	108(34)	31	34	30	72	75
	College and above	69(22)	29	25	23	59	72
χ^2 (df)			15.168(3)	10.300(3)	10.818(3)	12.576(3)	2.207(3)
p-value			.002*	.016*	.013*	.006*	.531
Depressive symptoms	No Depression	122(38)	15	11	9	59	67
	Depression	196(62)	56	51	48	83	81
χ^2 (df)			52.332(1)	51.132(1)	52.670(1)	21.613(1)	7.928(1)
p-value			.000*	.000*	.000*	.000*	.005*
Residence	Urban	234(74)	38	35	32	71	74
	Rural	39(12)	64	54	54	87	85
	Peri-Urban	45(14)	29	27	22	73	80
χ^2 (df)			12.138(2)	7.297(2)	10.058(2)	4.302(2)	2.756(2)
p-value			.002*	.026*	.007*	.116	.252
Systolic BP (n = 284)	Normal SBP	47(17)	28	23	19	60	66
	Prehypertension	189(67)	36	35	32	74	74
	Stage 1 HTN	29(10)	59	59	62	76	83
	Stage 2 HTN	19(7)	63	37	32	79	74
χ^2 (df)			12.643(3)	9.717(3)	15.200(3)	4.472(3)	2.623(3)
p-value			.005*	.021*	.002*	.215	.453

Results were expressed as %; Fisher's exact test was conducted when cells have expected count < 5

* p ≤ 0.05 was used as the limit for significance

Table 6 Correlation between selected independent variable with EQ-5D index and VAS

Dependent variables	EQ-5D index value		EQ-5D VAS	
	rho	^a p value	rho	^a p value
Age	-.212**	.000	.042	.459
Sex	-.271**	.000	-.157**	.005
Monthly family expenditure (BDT)	.072	.199	.220**	.000
Monthly family income (BDT)	.103	.067	.234**	.000
Duration of Diagnosis	-.122*	.030	-.011	.847
PHQ 9 score	-.616**	.000	-.380**	.000
BMI (Kg/ m ²)	.001	.986	.155*	.012
Education	.261**	.000	.205**	.000

^a P values were generated using Spearman's correlation coefficient

* Correlation is significant at the 0.05 level (2-tailed)

** Correlation is significant at the 0.01 level (2-tailed)

relationship between the severity of depression and HRQOL was significant at the 5% confidence level (Kruskal-Wallis Test statistics = 114.84; $p < .05$).

Discussion

This study examined the relationship between socio-demographic and anthropometric factors, comorbid depression, and HRQOL in adults with T2DM attending tertiary care hospitals in Dhaka, Bangladesh. The prevalence of depression and T2DM comorbid condition was 62%, which is higher than previous findings by Roy et al. [8] (34%) and Moussavi et al. [22] (9.3%). A meta-analysis by Anderson et al. [11] reported prevalence of depression ranged from 0.8 to 61% in T2DM patients. Studies in the USA and UK reported rates between 30 and 83% [54, 55], and an Iranian study found a 55% prevalence [56]. Studies from other countries indicated the prevalence of depression among T2DM patients of 37.6% in Saudi Arabia [57], 49.2% in Pakistan [25], and 17% in Dubai [58].

The higher prevalence of depression and T2DM comorbid conditions in this study may be attributed to two factors. Firstly, there was a higher proportion of women (58%) in our study compared to a previous study by Roy et al. (49%) [8]. This figure aligns with the findings from other studies that also indicated a high prevalence of depression among women [59, 60]. Second, we found a positive correlation between urban residency and depressive comorbidity. The current study had a higher proportion of urban residents (74%) than the previous study by Roy et al. (44.6%), potentially contributing to the increased prevalence of depression.

The findings of the EQ-5D in our study align with a study by Saleh et al., where most participants reported

experiencing anxiety/depression, while problems in self-care and usual activities were less reported [36]. However, the proportion of patients reporting problems in EQ-5D dimensions varies across countries. A study in Indonesia found lower proportions of problems in T2DM patients without comorbid depression (mobility 37%, self-care 12%, usual activities 23%, pain/discomfort 61%, and anxiety/depression 34%) compared to those with comorbid depression (mobility 36%, self-care 20%, usual activities 36%, pain/discomfort 69%, and anxiety/depression 44%) [61]. A population-based study in China reported lower problems in EQ-5D dimensions among people diagnosed with a chronic disease in the past six months (mobility 13%, self-care 8.2%, usual activities 12.4%, pain/discomfort 28%, and anxiety/depression 15%) [62]. These variations may be attributed to differences in population health status and demographic characteristics.

Depression has a negative impact on HRQOL, particularly in patients with comorbid diabetes [63]. Studies by Wallace et al. [64], Jing et al. [65], and Cannon et al. [21] have shown that depression in T2DM patients is associated with a poorer quality of life. Our study also found a significant association between depression and all five dimensions of EQ-5D in diabetic patients. Consistent with O'Neil et al. [66], we observed that the severity of depression correlates with changes in the health state of diabetic patients. Additionally, HRQOL declines with age, potentially due to the longer duration of illness, increased mental stress, and associated complications [25, 67, 68].

The current study revealed gender differences in HRQOL, with women reporting lower HRQOL. This is consistent with the gender differences in HRQOL observed in the general population [69, 70] and among people with diabetes [57, 61]. The female gender, along with hormonal and physiological characteristics, increases the risk of developing depression and subsequently lowers HRQOL [61, 71–73]. Consistent with previous studies [64, 65, 74], HRQOL deteriorates with a longer duration of diabetes, reflecting poor glycemic control, treatment compliance [65], and the development of diabetes-associated complications over the period [75]. It is evident from previous studies that laborers score low in HRQOL compared to other occupations, mainly due to stress related to work, financial adversity, and family problems [76, 77]. In contrast, our study found higher HRQOL scores among laborers, contrasting findings from other occupations. The reasons for this discrepancy require further investigation.

The study had some limitations, including the use of purposive and convenient sampling, a shorter data collection period, and the use of instruments to assess depression and HRQOL that are not considered the gold

Table 7 Mean PHQ 9 score, EQ-5D index and VAS by demographic characteristics (N= 318)

		N (%)	EQ-5D index Mean (SD)	EQ-5D VAS Mean (SD)	PHQ9 score Mean (SD)
Sex	Male	134(42)	.68(.19)	73(20)	5(4)
	Female	184(58)	.57(.23)	67(19)	9(5)
Age groups(years)	30–39	38(12)	.69(.21)	64(18)	7(5)
	40–49	79(25)	.64(.19)	70(21)	7(6)
	50–59	100(31)	.62(.19)	72(18)	7(5)
	60 +	101(32)	.56(.25)	68(21)	7(5)
Religion	Islam	312(98)	.61(.22)	69(20)	7(5)
	Hindu	6(2)	.64(.15)	73(17)	6(4)
Marital status	Never married	7(2)	.67(.25)	71(28)	10(8)
	Married	303(95)	.61(.22)	69(20)	7(5)
	Widow/Widower	7(2)	.69(.12)	79(11)	6(2)
	Separated/Divorced	1(1)	.72	70	9
Family History of T2DM	No	139(44)	.64(.18)	68(19)	7(5)
	Yes	179(56)	.60(.24)	70(20)	7(5)
Household members	≤ 4	145(46)	.64(.21)	70(20)	7(5)
	> 4	173(54)	.60(.22)	69(20)	8(5)
Education	No formal education	38(12)	.54(.26)	61(21)	10(6)
	Primary education	103(32)	.57(.24)	66(21)	8(5)
	Higher sec. education	108(34)	.64(.18)	72(18)	7(5)
	College and above	69(22)	.70(.18)	75(20)	5(4)
Occupation	Unemployed	3(1)	.56(.06)	65(22)	7(5)
	Service	54(17)	.68(.20)	74(23)	5(5)
	Business	32(10)	.66(.19)	72(23)	5(4)
	Labour	7(2)	.75(.22)	74(11)	8(6)
	Farming	5(2)	.63(.15)	60(10)	8(4)
	House wives	176(55)	.56(.22)	67(19)	9(5)
	Retired	41(13)	.70(.17)	74(16)	4(3)
Monthly household income(BDT)	< 10,000	52(16)	.53(.26)	63(20)	10(6)
	10,000–30000	146(46)	.63(.21)	67(20)	7(5)
	30,000–60000	83(26)	.62(.20)	73(18)	6(5)
	60,000–90000	11(3)	.69(.17)	81(14)	5(3)
	> 90,000	26(8)	.64(.19)	79(16)	6(4)
Monthly household expenditure(BDT)	< 10,000	58(18)	.56(.25)	64(19)	9(6)
	10,000–30000	155(49)	.63(.21)	68(20)	7(5)
	30,000–60000	80(25)	.62(.21)	73(19)	6(5)
	60,000–90000	9(3)	.72(.15)	83(16)	6(3)
	> 90,000	16(5)	.62(.21)	80(18)	7(4)
BMI	< 18.5 (Underweight)	9(3)	.62(.20)	66(20)	7(6)
	18.5 to < 23 (Normal wt.)	83(26)	.62(.20)	72(20)	7(5)
	23 to < 27.5 (Over wt.)	144(45)	.69(.20)	74(18)	6(6)
	≥ 27.5 (Obese)	82(26)	.55(.27)	69(27)	8(5)
Duration of diagnosis	Duration < = 3 years	108(34)	.66(.19)	70(18)	7(5)
	Duration 3–6 years	53(17)	.62(.20)	69(21)	7(5)
	Duration 6–9 years	47(14)	.59(.18)	69(18)	7(5)
	Duration > 9 years	110(35)	.58(.25)	69(22)	8(5)
Depressive symptoms	No depression(< 5 score)	122(38)	.75(.14)	76(16)	2(1)
	Depression(≥ 5 score)	196(62)	.53(.22)	65(21)	10(4)

Table 7 (continued)

		N (%)	EQ-5D index Mean (SD)	EQ-5D VAS Mean (SD)	PHQ9 score Mean (SD)
Residence	Urban	234(74)	.63(.21)	71(20)	7(5)
	Rural	39(12)	.49(.27)	60(21)	10(6)
Systolic BP	Peri-Urban	45(14)	.67(.15)	70(18)	6(4)
	Normal SBP	47(17)	.72(.16)	68(23)	7(4)
	Prehypertension	189(67)	.63(.20)	71(18)	6(5)
	Stage 1 Hypertension	29(10)	.49(.25)	67(25)	10(5)
	Stage 2 Hypertension	19(7)	.57(.22)	66(19)	10(6)




Fig. 3 Mean EQ-5D index and severity of depression: displays the relationship between the mean EQ-5D-5L index score, a measure of HRQOL, and the severity of depression among the study population. The x-axis represents the severity of depression, measured using a validated questionnaire, and is divided into four categories: no depression, mild depression, moderate depression, and severe depression. The y-axis shows the mean EQ-5D-5L index score for each category of depression severity. The legend provides information about the mean EQ-5D-5L index score for each category of depression severity, allowing readers to interpret the relationship between depression severity and HRQOL in the study population. The graph shows a clear decrease in the mean EQ-5D-5L index score as depression severity increases, indicating that depression is associated with lower HRQOL.

standards. The PHQ-9 and EQ-5D rely on self-reported data and may be less accurate and sensitive for certain populations. The gold standard for assessing depression is typically a structured clinical interview using standardized diagnostic criteria, such as DSM-5 or ICD-11, conducted by trained mental health professionals. The study was cross-sectional with small samples, limiting causal or dose-response conclusions. Generalizability is restricted,

as most patients were females from two hospitals in Dhaka, Bangladesh.

Conclusions

A significant proportion of patients with T2DM in this study reported having depression and T2DM comorbid conditions. Further, depressive symptoms negatively impact the HRQOL, with the patients experiencing lower

HRQOL as the severity of depression increases. These findings provide an important basis for considering early screening, diagnosis, treatment, and follow-up of depression in people with diabetes. By identifying an association between independent socio-demographic variables, depressive comorbidity, and HRQOL, this study offers scope for designing and conducting further research at a larger scale employing more rigor.

Abbreviations

BIHS	Bangladesh Institute of Health Science
BRAC	Bangladesh Rural Advancement Committee
BMI	Body Mass Index
CIDI	Composite International Diagnostic Interview
DMCH	Dhaka Medical College and Hospital
EuroQoL-5D-5L	European Quality of Life Five Dimension and Five Level
HCDP	Health Care Development Project
HRQOL	Health-Related Quality of Life
IDF	International Diabetes Federation
NCD	Non-Communicable disease
OPD	Outpatient Department
PHQ-9	Patient Health Questionnaire (PHQ)-9
QoL	Quality of Life
SPSS	Statistical Package for Social Sciences
T2DM	Type 2 diabetes mellitus
UHC	Urban Health Centers
VAS	Visual Analogue Scale

Acknowledgements

We sincerely thank to Ms. Nadia Alamgir and Mr. Kuhel Islam for their guidance and Dr. Farhana Karim, Mr. Pradeep Adhikari, Dr. Punam Ganguly, Mr. Kaoser Bin Siddique, Mr. Mahfuz Rahman for their support while conducting this study. The head of the department- Endocrinology, Dhaka Medical College, and Hospital, Superintendent and Director, Bangladesh Institute of Health Sciences, for permission to collect the data in the institute premises. We also would like to thank the academic and management team of JPGSPH, Dhaka, for their support and guidance throughout this study. Further, we are grateful to Mr. Hassan Benya (MPH) from Freetown-Sierra Leone for his English proof-read of this paper. Lastly, we convey sincere gratitude to all the participants who participated in the study and provided valuable information.

Author contributions

Study conception and design, data collection: MKN; analysis and interpretation of results: MKN, LBR, SV; draft manuscript preparation: MKN, SV, RI, SN, RDG. All authors (MKN, SV, RI, SN, RDG, LBR) reviewed the results and approved the final version of the manuscript.

Funding

Not applicable.

Availability of data and materials

All data generated or analyzed during this study are included in this published article as annexed tables.

Declarations

Ethics approval and consent to participate

Ethical approval was obtained from the institutional review committee of the JP Grant School of Public Health, BRAC (Bangladesh Rural Advancement Committee) University. Prior permission to conduct the study was sought from BIHS and DMCH authorities. Informed written consent was obtained from all the patients after fully explaining the nature, purpose, and procedures used for the study. Participants were informed about the confidentiality of the information, and the ID numbers of patients were used to maintain anonymity. The right to withdraw from the interview at any time and future benefits of the study were also communicated to participants.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Independent Scholar and Alumni, JPGSPH, BRAC University, Bangladesh, Chhindwara, India. ²Tata Institute of Social Sciences, Mumbai, India. ³University of South Carolina, Columbia, SC, USA. ⁴International Initiative for Impact Evaluation (3ie), Columbia, SC, USA. ⁵University of Texas at El Paso, El Paso, TX, USA. ⁶Central Queensland University, Sydney Campus, Sydney, Australia.

Received: 14 September 2022 Accepted: 3 October 2023

Published online: 16 October 2023

References

- International Diabetes Federation. IDF Diabetes Atlas. 10th ed. Brussels, Belgium: International Diabetes Federation; 2021.
- International Diabetes Federation. IDF Diabetes Atlas. vol. 266. 9th ed. Brussels, Belgium: International Diabetes Federation; 2019.
- Bhowmik B, Afsana F, My Diep L, Munir SB, Wright E, Mahmood S, et al. Increasing prevalence of type 2 diabetes in a rural Bangladeshi population: a population based study for 10 years. *Diabetes Metab J*. 2013;37:46–53. <https://doi.org/10.4093/dmj.2013.37.1.46>.
- Global Health Data Exchange. GBD Results Tool|GHDx. Institute for Health Metrics and Evaluation 2019.
- Biswas T, Islam MS, Linton N, Rawal LB. Socio-economic inequality of chronic non-communicable diseases in Bangladesh. *PLoS ONE*. 2016. <https://doi.org/10.1371/journal.pone.0167140>.
- Lahiry S. Cost of Diabetes care in the Out-patient Department of A Tertiary Care Hospital in Bangladesh. Oslo: University of Oslo; 2006.
- Pan A, Lucas M, Sun Q, Van Dam RM, Franco OH, Manson JE, et al. Bidirectional Association Between Depression and Type 2 Diabetes Mellitus in Women. n.d.
- Roy T, Lloyd CE, Parvin M, Galib K, Mohiuddin B, Rahman M. Prevalence of co-morbid depression in out-patients with type 2 diabetes mellitus in Bangladesh. *Science*. 2012;6:98.
- Hosen I, Al-Mamun F, Mamun MA. Prevalence and risk factors of the symptoms of depression, anxiety, and stress during the COVID-19 pandemic in Bangladesh: a systematic review and meta-analysis. *Global Mental Health*. 2021;8:e47. <https://doi.org/10.1017/gmh.2021.49>.
- Dejean D, Giacomini M, Vanstone M, Brundisini F. Patient Experiences of depression and anxiety with chronic disease: a systematic review and qualitative meta-synthesis. *Science*. 2013;13:46.
- Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care*. 2001. <https://doi.org/10.2337/diacare.24.6.1069>.
- Musselman DL, Betan E, Larsen H, Phillips LS. Relationship of depression to diabetes types 1 and 2: epidemiology, biology, and treatment. *Biol Psychiatry*. 2003. [https://doi.org/10.1016/S0006-3223\(03\)00569-9](https://doi.org/10.1016/S0006-3223(03)00569-9).
- Alzoubi A, Abunaser R, Khassawneh A, Alfaqih M, Khasawneh A, Abdo N. The bidirectional relationship between diabetes and depression: a literature review. *Korean J Fam Med*. 2018;39:137–46. <https://doi.org/10.4082/kjfm.2018.39.3.137>.
- Talbot F, Nouwen A. A review of the relationship between depression and diabetes in adults: is there a link? *Diabetes Care*. 2000;23:1556–62. <https://doi.org/10.2337/DIACARE.23.10.1556>.
- Zhao F, Pan A, Yang X, Meng R, Ye Y, Wang Y, et al. Bidirectional association between depressive symptoms and type 2 diabetes mellitus: the China Health and Retirement Longitudinal Study. *J Diabetes Complications*. 2019;33:107387. <https://doi.org/10.1016/j.jdiacomp.2019.05.018>.
- Chamberlain AM, McNallan SM, Dunlay SM, Spertus JA, Redfield BM, Moser DK, et al. Physical health status measures predict all-cause mortality in patients with heart failure. *Circ Heart Fail*. 2013. <https://doi.org/10.1161/CIRCHEARTFAILURE.112.000291>.
- Irfan SN, Rahaman MM, Noman A, Mithun S. Health related quality of life among coronary artery bypass graft patient attended at combined

- military Hospital, Dhaka. Anwer Khan Modern Medical College J. 2013. <https://doi.org/10.3329/akmmcj.v4i2.16918>.
18. Carr AJ, Gibson B, Robinson PG. Measuring quality of life is quality of life determined by expectations or experience? *Br Med J*. 2001. <https://doi.org/10.1136/bmj.322.7296.1240>.
19. Centers for Disease Control and Prevention. HRQOL Concepts. Centers for Disease Control and Prevention 2016.
20. Bonomi AE, Patrick DL, Bushnell DM, Martin M. Validation of the United States' version of the World Health Organization Quality of Life (WHO-QOL) instrument. *J Clin Epidemiol*. 2000. [https://doi.org/10.1016/S0895-4356\(99\)00123-7](https://doi.org/10.1016/S0895-4356(99)00123-7).
21. Cannon A, Handelsman Y, Heile M, Shannon M. Burden of Illness in Type 2 diabetes mellitus. *J Manag Care Spec Pharm*. 2018;5:24.
22. Moussavi S, Chatterji S, Verdes E, Tandon A, Patel V, Ustun B. Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *Lancet*. 2007. [https://doi.org/10.1016/S0140-6736\(07\)61415-9](https://doi.org/10.1016/S0140-6736(07)61415-9).
23. Otley A, Smith C, Nicholas D, Munk M, Avolio J, Sherman PM, et al. The IMPACT questionnaire: a valid measure of health-related quality of life in pediatric inflammatory bowel disease. *J Pediatr Gastroenterol Nutr*. 2002. <https://doi.org/10.1097/00005176-200210000-00018>.
24. O'Neil A, Taylor B, Sanderson K, Cyril S, Chan B, Hawkes AL, et al. Efficacy and feasibility of a tele-health intervention for acute coronary syndrome patients with depression: results of the "MoodCare" randomized controlled trial. *Ann Behav Med*. 2014. <https://doi.org/10.1007/s12160-014-9592-0>.
25. Khan P, Qayyum N, Malik F, Khan T, Khan M, Tahir A. Incidence of anxiety and depression among patients with type 2 diabetes and the predicting factors. *Cureus*. 2019. <https://doi.org/10.7759/cureus.4254>.
26. Baumeister H, Balke K, Härter M. Psychiatric and somatic comorbidities are negatively associated with quality of life in physically ill patients. *J Clin Epidemiol*. 2005. <https://doi.org/10.1016/j.jclinepi.2005.03.011>.
27. Martínez YV, Prado-Aguilar CA, Rascón-Pacheco RA, Valdivia-Martínez JJ. Quality of life associated with treatment adherence in patients with type 2 diabetes: a cross-sectional study. *BMC Health Serv Res*. 2008. <https://doi.org/10.1186/1472-6963-8-164>.
28. Rejeski WJ, Wei L, Neiberg RH, van Dorsten B, Foster GD, Maciejewski ML, et al. Correlates of health-related quality of life in overweight and obese adults with type 2 diabetes. *Obesity*. 2006. <https://doi.org/10.1038/oby.2006.101>.
29. Vilhena E, Pais-Ribeiro J, Silva I, Pedro L, Meneses RF, Cardoso H, et al. Psychosocial factors as predictors of quality of life in chronic portuguese patients. *Health Qual Life Outcomes*. 2014. <https://doi.org/10.1186/1477-7525-12-3>.
30. Katon WJ, Lin EHB, von Korff M, Ciechanowski P, Ludman EJ, Young B, et al. Collaborative care for patients with depression and chronic illnesses. *New England J Med*. 2010. <https://doi.org/10.1056/nejmoa1003955>.
31. Chew BH, Vos RC, Metzendorf MI, Scholten RJPM, Rutten GEHM. Psychological interventions for diabetes-related distress in adults with type 2 diabetes mellitus. *Cochrane Database Syst Rev*. 2017. <https://doi.org/10.1002/14651858.CD011469.pub2>.
32. Power R, Galea C, Muhit M, Heanoy E, Karim T, Badawi N, et al. What predicts the proxy-reported health-related quality of life of adolescents with cerebral palsy in Bangladesh? *BMC Public Health*. 2020. <https://doi.org/10.1186/s12889-019-8130-1>.
33. Bowers B, Ruth Butlin C, Alam K, Lockwood DN, Walker SL, Health Programme R. Health-Related Quality of Life amongst people affected by Erythema Nodosum Leprosum in Bangladesh: a Cross Sectional Study. n.d.
34. Ahmmed FS, Rahman MS, Zafreen F, Ara R, Islam MZ. Health-related quality of life among elderly population of Bangladesh. *J Armed Forces Med College Bangladesh*. 2020;15:161–4. <https://doi.org/10.3329/jafmc.v15i2.50828>.
35. Mahumud RA, Ali N, Sheikh N, Akram R, Alam K, Gow J, et al. Measuring perinatal and postpartum quality of life of women and associated factors in semi-urban Bangladesh. *Qual Life Res*. 2019. <https://doi.org/10.1007/s11136-019-02247-0>.
36. Saleh F, Mumu SJ, Ara F, Hafez MA, Ali L. Non-adherence to self-care practices & medication and health related quality of life among patients with type 2 diabetes: a cross-sectional study. *BMC Public Health*. 2014. <https://doi.org/10.1186/1471-2458-14-431>.
37. Barua L, Faruque M, Chowdhury HA, Banik PC, Ali L. Health-related quality of life and its predictors among the type 2 diabetes population of Bangladesh: a nation-wide cross-sectional study. *J Diabetes Investig*. 2021;12:277–85. <https://doi.org/10.1111/jdi.13331>.
38. Islam M, Ahmed S, Sarker R, Farjana S, Akter A, Saha S. Health-related quality of life among adult migrant garment workers in Dhaka City. *Bangladesh Med J*. 2014. <https://doi.org/10.3329/bmj.v40i3.18651>.
39. Alessi J, de Oliveira GB, Franco DW, Brino DAB, Becker AS, Knijnik CP, et al. Mental health in the era of COVID-19: Prevalence of psychiatric disorders in a cohort of patients with type 1 and type 2 diabetes during the social distancing. *Diabetol Metab Syndr*. 2020. <https://doi.org/10.1186/s13098-020-00584-6>.
40. Editor T. Contribution of Bangladesh Diabetic Samity (BADAS) to Health Care Management. *BIRDEM Med J* 2012; 2. <https://doi.org/10.3329/birdem.v2i1.12352>.
41. Dhaka Medical College. Dhaka Medical College. 2021 n.d. <http://dmc.gov.bd/site/page/1cb15259-e127-4705-bb31-116f5dd48052/> (Accessed 7 Oct 2021).
42. da Costa D, Dritsa M, Verreault N, Balaa C, Kudzman J, Khalifé S. Sleep problems and depressed mood negatively impact health-related quality of life during pregnancy. *Arch Womens Ment Health*. 2010. <https://doi.org/10.1007/s00737-009-0104-3>.
43. Roset M, Badia X, Mayo NE. Sample size calculations in studies using the EuroQol 5D. *Qual Life Res*. 1999. <https://doi.org/10.1023/A:1008973731515>.
44. Dean AG, Sullivan KM, Soe MM. OpenEpi: Open Source Statistics for Public Health, Version 3.01. OpenEpi 2013. <https://www.openepi.com/>.
45. Heyworth ITM, Hazell ML, Linehan MF, Frank TL. How do common chronic conditions affect health-related quality of life? *Br J Gen Pract*. 2009. <https://doi.org/10.3399/bjgp09X453990>.
46. Herdman M, Gudex C, Lloyd A, Janssen M, Kind P, Parkin D, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res*. 2011. <https://doi.org/10.1007/s11136-011-9903-x>.
47. van Hout B, Janssen MF, Feng YS, Kohlmann T, Busschbach J, Golicki D, et al. Interim scoring for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets. *Value Health*. 2012. <https://doi.org/10.1016/j.jval.2012.02.008>.
48. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. 2001. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x>.
49. Kochhar P, Rajadhyaksha S, Suvarna V. Translation and validation of brief patient health questionnaire against DSM IV as a tool to diagnose major depressive disorder in Indian patients. *J Postgrad Med*. 2007. <https://doi.org/10.4103/0022-3859.32209>.
50. Kroenke K, Spitzer RL. The PHQ-9: a new depression diagnostic and severity measure. *Psychiatr Ann*. 2002. <https://doi.org/10.3928/0048-5713-20020901-06>.
51. IBM Corp. Released. IBM SPSS Statistics for Windows, Version 20.0. 2011.
52. Hazell M, Frank T, Frank P. Health related quality of life in individuals with asthma related symptoms. *Respir Med*. 2003. [https://doi.org/10.1016/S0954-6111\(03\)00249-X](https://doi.org/10.1016/S0954-6111(03)00249-X).
53. van Reenen M, Janssen B, Stolk E, Boye KS, Herdman M, Kennedy-Martin M, et al. CHANGES INCLUDED in this update of the EQ-5D-5L User Guide Table of contents. n.d.
54. Kendrick T, Dowrick C, McBride A, Howe A, Clarke P, Maisey S, et al. Management of depression in UK general practice in relation to scores on depression severity questionnaires: analysis of medical record data. *BMJ (Online)*. 2009. <https://doi.org/10.1136/bmj.b750>.
55. Li C, Ford ES, Strine TW, Mokdad AH. Prevalence of depression among U.S. adults with diabetes: findings from the 2006 behavioral risk factor surveillance system. *Diabetes Care*. 2008. <https://doi.org/10.2337/dc07-1154>.
56. Khamseh ME, Baradaran HR, Rajabali H. Depression and diabetes in Iranian patients: a comparative study. *Int J Psychiatry Med*. 2007. <https://doi.org/10.2190/FP64-82V3-1741-842V>.
57. Albasheer OB, Mahfouz MS, Solan Y, Khan DA, Muqri MA, Almutairi HA, et al. Depression and related risk factors among patients with type 2 diabetes mellitus, Jazan area, KSA: a cross-sectional study. *Diabetes Metab Syndr*. 2018;12:117–21. <https://doi.org/10.1016/j.dsx.2017.09.014>.
58. Alajmani DSA, Alkaabi AM, Alhosani MW, Foad AA, Abdouli FA, Carrick FR, et al. Prevalence of undiagnosed depression in patients with type 2

- diabetes. *Front Endocrinol* (Lausanne). 2019. <https://doi.org/10.3389/fendo.2019.00259>.
59. Ali S, Stone MA, Peters JL, Davies MJ, Khunti K. The prevalence of co-morbid depression in adults with Type 2 diabetes: a systematic review and meta-analysis. *Diabetic Med*. 2006. <https://doi.org/10.1111/j.1464-5491.2006.01943.x>.
60. Asghar S, Hussain A, Ali SMK, Khan AKA, Magnusson A. Prevalence of depression and diabetes: a population-based study from rural Bangladesh. *Diabetic Med*. 2007. <https://doi.org/10.1111/j.1464-5491.2007.02136.x>.
61. Arifin B, Idrus LR, van Asselt ADI, Purba FD, Perwitasari DA, Thobari JA, et al. Health-related quality of life in Indonesian type 2 diabetes mellitus outpatients measured with the Bahasa version of EQ-5D. *Qual Life Res*. 2019;28:1179–90. <https://doi.org/10.1007/s11136-019-02105-z>.
62. Sun S, Chen J, Johannesson M, Kind P, Xu L, Zhang Y, et al. Population health status in China: EQ-5D results, by age, sex and socio-economic status, from the national health services survey 2008. *Qual Life Res*. 2011. <https://doi.org/10.1007/s11136-010-9762-x>.
63. Liu X, Haagsma J, Sijbrands E, Buijks H, Boogaard L, Mackenbach JP, et al. Anxiety and depression in diabetes care: longitudinal associations with health-related quality of life. *Sci Rep*. 2020. <https://doi.org/10.1038/s41598-020-57647-x>.
64. Wallace K, Zhao X, Misra R, Sambamoorthi U. The humanistic and economic burden associated with anxiety and depression among adults with comorbid diabetes and hypertension. *J Diabetes Res*. 2018. <https://doi.org/10.1155/2018/4842520>.
65. Jing X, Chen J, Dong Y, Han D, Zhao H, Wang X, et al. Related factors of quality of life of type 2 diabetes patients: a systematic review and meta-analysis 11 Medical and Health Sciences 1117 Public Health and Health Services 11 Medical and Health Sciences 1103 Clinical Sciences. *Health Qual Life Outcomes*. 2018. <https://doi.org/10.1186/s12955-018-1021-9>.
66. O'Neil A, Stevenson CE, Williams ED, Mortimer D, Oldenburg B, Sanderson K. The health-related quality of life burden of co-morbid cardiovascular disease and major depressive disorder in Australia: findings from a population-based, cross-sectional study. *Qual Life Res*. 2013. <https://doi.org/10.1007/s11136-012-0128-4>.
67. Nilsson J, Rana AKMM, Kabir ZN. Social capital and quality of life in old age: results from a cross-sectional study in rural Bangladesh. *J Aging Health*. 2006. <https://doi.org/10.1177/0898264306286198>.
68. Sazlina SG, Zaiton A, Nor Afiah MZ, Hayati KS. Predictors of health related quality of life in older people with non-communicable diseases attending three primary care clinics in Malaysia. *J Nutr Health Aging*. 2012. <https://doi.org/10.1007/s12603-012-0038-8>.
69. Malfa CS, Karaivazoglou K, Assimakopoulos K, Gourzis P, Vantarakis A. Psychological distress and health-related quality of life in public sector personnel. *Int J Environ Res Public Health*. 2021. <https://doi.org/10.3390/ijerph18041865>.
70. del Río LM, García-Calvente MM, Calle-Romero J, Machón-Sobrado M, Larrañaga-Padilla I. Health-related quality of life in Spanish informal caregivers: gender differences and support received. *Qual Life Res*. 2017. <https://doi.org/10.1007/s11136-017-1678-2>.
71. Astbury J. Gender and mental health. Working Paper Series No. 99. Cambridge, MA: Harvard Center for Population and Development Studies; 1999. p. 8.
72. Kristensen Maria. Health related quality of life and compliance in rheumatoid arthritis patients treated with biologic subcutaneous drugs: A cross sectional study. Unpublished dissertation. Aalborg University, 2012.
73. Zelinkova Z. On reproduction and other gender and sex-related issues in inflammatory bowel disease (Inflammatory bowel disease du deuxième-sexe). 2012.
74. Romero-Naranjo F, Espinosa-Uquillas C, Gordillo-Altamirano F, Barrera-Guarderas F. Which factors may reduce the health-related quality of life of ecuadorian patients with diabetes? *P R Health Sci J*. 2019;38:102–8. <https://doi.org/10.1097/01.jom.0000052965.43131.c3>.
75. Trikkalinou A, Papazafropoulou AK, Melidonis A. Type 2 diabetes and quality of life. *World J Diabetes*. 2017;8:120. <https://doi.org/10.4239/wjcd.v8.i4.120>.
76. Riise T, Moen BE, Nortvedt MW. Occupation, lifestyle factors and health-related quality of life: the Hordaland Health Study. *J Occup Environ Med*. 2003. <https://doi.org/10.1097/01.jom.0000052965.43131.c3>.
77. Gebru T, Lentiro K. The impact of community-based health insurance on health-related quality of life and associated factors in Ethiopia: a comparative cross-sectional study. *Health Qual Life Outcomes*. 2018. <https://doi.org/10.1186/s12955-018-0946-3>.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

